

EFFECTS OF CHLORINE DIOXIDE AND SOME WATER QUALITY PARAMETERS ON THE FORMATION OF THMS IN WATER TREATMENT PLANTS

ALAAHUSAEEN AL-FATLAWI

Assistant Professor, Babylon University, College of Engineering, Babylon, Iraq

ABSTRACT

The presence of trihalomethanes (THMs) in drinking water has attracted the attention of both researchers and professionals, because of the harmful effects of these substances on human health. Chlorine dioxide (ClO_2) is an alternative to chlorine because it is an oxidizing agent rather than a chlorinating agent, and therefore, will not form chlorinated disinfection byproducts such as THMs under typical water treatment conditions.

The present study was conducted to determine the effects of Chlorine Dioxide, Temperature, Turbidity, Hydrogen ion concentration (pH), Total Organic Carbon (TOC), and contact time on the formation of THMs during treatment process. A small-scale pilot plant of a compact model is designed and constructed at Environmental Engineering Department lab (Babylon University). It is simple in operate and made of galvanized iron of its components and locally available materials.

Total Trihalomethanes (TTHMs), their speciation compounds (THM4), and varieties of water quality parameters were monitored by addition different doses of chlorine dioxide and alum to raw water pipe entering the pilot plant. The input flow rate was ($0.36 \text{ m}^3/\text{hr}$) with characteristics (as average) (Temperature, Turbidity, Alkalinity, TTHMs, pH, and TOC) are (28 $^\circ\text{C}$, 27 NTU, 140 mg/L, 0.1545 mg/L, 8.22, 1.4 mg/L) respectively.

All runs were done during April, May and June of 2012. The linear correlations were calculated between THM concentrations and ClO_2 dose or selected water quality parameters, in an attempt to identify one or more parameters that could be used as surrogate in THMs investigation. (HS – GC - ECD) with Gas chromatography analysis techniques were used to measure the THMs concentrations. It was noticed that TTHMs concentration increases as temperature, TOC, and pH increase, and average TTHM levels detected in all runs, does not exceed the USEPA's Stage I (limit of 80 $\mu\text{g/L}$), while exceeded the stage II limit in all sampling measurements.

KEYWORDS: Trihalomethane, Drinking Water, Temperature, Turbidity, pH, TOC

INTRODUCTION

In 1976 the U.S. National Cancer Institute published a report linking chloroform to cancer in laboratory animals. Two years previously, (Rook, 1974), was the first to report on the occurrence of this compound in chlorinated drinking water. During the ensuing years water with disinfectant chemical reagents, as a result of which a long and varied list of disinfection by-products (DBPs)--many of which might be teratogenic, mutagenic and/or carcinogenic--has been compiled (Fawell and Hunt, 1988) in which the trihalomethanes (THMs) are only the tip of the iceberg (Klein, 1990). The concern of the national and supranational health authorities has resulted either in recommendations or regulations about the maximum levels of several groups of DBPs (EEC, 1980; USEPA, 1988; WHO, 1992) and has prompted studies

on the reaction mechanisms and conditions involved and, furthermore, on developing kinetic computer models to simulate the formation of DBPs during water treatment, for both their control and minimization. The information on the reaction mechanism of the formation of THMs is still limited, although it is generally recognized that four factors would be involved in their formation: the chlorine-to-precursor molar ratio, pH, temperature and reaction time.

THM formation is strongly dependent upon the chlorine concentration (Kavanough et al., 1980; Peters et al., 1980). However, there is some disagreement regarding the quantitative relations between chlorine concentrations and the rate of THMs production, linear relationship between chlorine consumption and the production of THMs with a reaction order greater or equal to unity (Kavanough et al., 1980; Trussel and Umphres, 1978). Despite this, it is also possible that the reaction order might change during the course of the reaction (Kavanough et al., 1980).

The formation of THMs also increases strongly with increasing amounts of soluble organic matter, following a first-order reaction. In naturally occurring water, this organic matter usually consists of humic substances (Trussel and Umphres, 1978; Babcock and Singer, 1979). Although fulvic acid accounts for over 90% of the aqueous humics in many water sources, Babcock and Singer found that relative contributions to the formation of THMs by the humic fraction is greater than that of the fulvic fraction since the former substances react more readily with chlorine.

Increased pH values lead to increases in THMs formation (Stevens et al., 1989; Onodera et al., 1987), three-fold increases being reported in the reaction rate per unit of pH (Kavanough et al., 1980). The lower the pH, the higher non-ionized HClO form of hypochlorous acid is found, thus increasing its reaction rate with the humic matter. However, THM yields depend rather on the last step of the THM reaction pathway, which is base-catalyzed as with the haloform reaction (Simmon and Tardiff, 1978). These findings have also been reported by other authors (Peters et al., 1980; Sandier, 1977). According to Abedin et al., (2010), the acidic functional groups of humic matter are not ionized, leading to the aggregation of molecules due to Van der Waals forces. This phenomenon is also associated with folding of the humic molecules, leaving fewer sites available for attack by chlorine (Trehy and Bieber, 1980), thereby reducing THM production. In studies on the effect of temperature on THMs formation, an Arrhenius-type dependence has been found between the rate constant and temperature, with activation energies ranging from 10-20 kJ mol (Peters et al., 1980) to below 10 kJ mol (Kavanough et al., 1980; Stevens et al., 1989). Accordingly, a higher rate of THM formation should be expected with increasing water temperatures although, on the other hand, the volatility of these compounds should account for their partial removal in open systems.

EXPERIMENTAL WORK

This paper deals with the analyses and discussion of the results of experimental work to show the effect of ClO₂ on the formation of THMs and other water quality parameters. Total Trihalomethanes (TTHMs), their speciation compounds (THM4), and varieties of water quality parameters were monitored in a small-scale pilot plant. By addition different doses of chlorine dioxide and alum to raw water pipe entering the pilot plant and monitoring THMs concentration. The raw and finished water quality parameters which were analyzed are included; Hydrogen ion concentration, Temperature, Total organic carbon, Turbidity, Total Trihalomethane concentration and examines the percent reduction of Trihalomethanes formation with applying different doses of chlorine dioxide. All runs were done during April, May and June of 2012. The linear correlations were calculated between THM concentrations and ClO₂ dose or selected water quality parameters, in an attempt to identify one or more parameters that could be used as surrogate in THMs investigation. The input flow rate was (0.36 m³/hr) with characteristics listed in Table (1):


Table 1: Characteristics of Raw Water

Parameter		Value, (Average)
Temperature, ($^{\circ}\text{C}$)		28
Turbidity, (NTU)		27
Alkalinity, (mg/L)		140
THMs (mg/L)	CF	0.02400
	BDCM	0.05257
	DCBM	0.0588
	BF	0.01905
TTHMs, (mg/L)		0.1545
pH		8.22
TOC, (mg/L)		1.4

TTHM SPECIATION IN THE STUDY

In this study another point of interest was the THM speciation. Figure 9 presents a representative THM speciation. This figure summarizes the weight fractions of CHCl_3 , CHBrCl_2 , CHBr_2Cl , and CHBr_3 species. Results show that the intermediate brominated species (CHBrCl_2 , and CHBr_2Cl) comprised most of the THM concentrations. However, results of recent toxicological studies suggest further attention should be paid to the occurrence of brominated compounds, especially bromodichloromethane (CHBrCl_2) (Krasner et al., 1992).

In general, all results indicate that the treated water contain mainly the brominate THM species (CHClBr_2 and CHBrCl_2). The average weight fraction of chloroform (CHCl_3) compound ranged from 7.16% to 21.46%, while the average weight fraction of the rest brominated (CHBr_2Cl , and CHBrCl_2) species ranged from about 34.97% to 43.8%, 30.67% to 34.54% respectively as shown in Figure 1. Nearly the same pattern was pointed out by (McLain et al., 2002).

**Figure 1: Weight Fraction of Individual THM4 to Total THM4**

EFFECTS OF CHLORINE DIOXIDE DOSE ON TRIHALOMETHAN FORMATION

The results showed that the THM formation decrease with increasing chlorine dioxide dose, concentrations of THMs that developed in water pretreated with 2 mg/L ClO_2 were significantly lower when compared to those in water pretreated with the other dose of ClO_2 . It is observed that higher THM level at ClO_2 dose of 0.2 mg/l and then tends to decrease slightly with increasing ClO_2 dose. Increasing ClO_2 doses have been shown to have the opposite effect as chlorine. This finding is agreed with the results adapted by (Griese, 1991).

Also the results showed that with increasing dose of ClO_2 from (1.0 to 2.0) mg/L leads to decrease reduction in THMs concentration with about (24) percent (Griese, 1991; Hulsey et al., 2000) were also found the same results. They were found that increasing doses of ClO_2 leads to increasing reduction in THM formation and approximately (20.74) percent reduction in THM concentrations when 1.0 to 2 mg/L of ClO_2 was applied.

Figure 2 shows the variation of TTHMs with chlorine dioxide dose used in the study. As shown the TTHMs formation decreasing with increasing chlorine dioxide dose. This Figure was performed on the ClO_2 samples to determine whether ClO_2 dose affected the final TTHM levels.

While Figure 3 represents the percent reduction in TTHMs concentration of filtered water after pre oxidation with ClO_2 . From this Figure we found that the removal efficiency increases with the increase chlorine dioxide dose, similar result was found by (Hulsey et al., 2000) who concluded that the removal efficiency of THMs from water increase by increasing chlorine dioxide dose.

As expected, correlations are negative between THM4 concentrations and chlorine dioxide dose. During study period, the regression equation indicates that higher percent removal of TTHMs was (64.33%) when using (2mg/L) of ClO_2 dose. The present results showed that the higher dose of ClO_2 is considerably efficient for the removal of THMs from raw water. The performance of the ClO_2 in reduction of THMs formation can be expressed in the form $Y=7.637X+46.76$ with coefficient of determination (R^2) equal to 0.902

Figure 3 shows that the THMs removal efficiency is highly dependent on chlorine dioxide dosage. It is evident that at higher dosages (more than 1.5 mg/L), the ClO_2 started to achieve higher removal percentages of THMs. This can be attributed to the fact that the chlorine dioxide has higher capacity to removal THMs.




Figure 2: The Variation of TTHMs with ClO_2 Dose




Figure 3: The Percent Reductions in the TTHMs Concentrations after ClO_2 Dose

Figure 4 presented a comparison of total THMs measurements with the USEPA Stage I and stage II limits. None of the THM levels exceeded the stage I limit, while, the stage II limit was exceeded in all sampling measurements.




Figure 4: Maximum THMs Levels Detected in the Study

Figure 5 shows the variation of four THMs compound with ClO₂ dose. As shown the dominant TTHMs species during all runs is DCBM, it consists (39.2%) of trihalomethanes. It decreases slightly with increasing dose, the remaining THMs are bromodichloromethane 33.34% which began slightly decrease with increasing dose, while the chloroform is 14.5%, and bromoform is 8.6%.




Figure 5: Relative Amounts of THMs Compounds with ClO₂ Dose

Figures 6 and 7 show the relation between chlorine dioxide dosing and percent reduction of two compounds (chloroform, and bromodichloromethane). Correlations are negative between chloroform percent reduction and chlorine dioxide dose under the dose (1 mg/L), above this dose the percent began to increase, so the chloroform percent reduction increases with increasing chlorine dioxide dose above (1 mg/L). Thus, we conclude that chloroform concentration is proportional with ClO₂ dose in the form ($Y = 5.3961X + 52.04$) with coefficient of determination ($R^2 = 0.0407$). While correlations are positive between bromodichloromethane percent reduction and chlorine dioxide dose (decrease in BDCM concentrations with increasing chlorine dioxide dose). Thus, we conclude that BDCM concentration is proportional with ClO₂ dose in the form ($Y = 8.846X + 46.408$) with coefficient of determination ($R^2 = 0.9122$) as shown in Figure 7.




Figure 6: Percent Reduction in CF with the ClO₂ Dose




Figure 7: Percent Reduction in BDCM with the ClO₂ Dose

The correlation is positive between dichlorobromomethane percent reduction and chlorine dioxide dose (decrease in DCBM concentrations with increasing chlorine dioxide dose). Thus, we conclude that DCBM concentration is proportional with ClO₂ dose in the form ($Y = 8.298x + 44.76$) with coefficient of determination ($R^2 = 0.752$) see Figure 8. While positive correlations between bromoform percent reduction and chlorine dioxide dose (slightly decrease in BF concentrations with increasing chlorine dioxide dose). Thus, we conclude that bromoform concentration is proportional with ClO₂ dose in the form ($Y = 4.927X + 49.65$) with weak coefficient of determination ($R^2 = 0.28$) as shown in Figure 9.




Figure 8: Percent Reduction in DCBM with the ClO₂ Dose




Figure 9: Percent Reduction in BF with the ClO₂ Dose

EFFECT OF CONTACT TIME ON TTHMS CONCENTRATION

The effect of detention time on TTHMs formation was investigated with a dosage of 2mg/L chlorine dioxide. The concentration of TTHMs was measured after 30, 60, 90 and 120 min. Figure 10 shows that the TTHMs concentration decreases significantly with the increasing contact time. Thus, we conclude that the THMs concentration is proportional with time in the form ($Y = -0.0009X + 0.1088$) with coefficient of determination $R^2 = 0.88$. The increasing contact time of ClO₂ leads to decrease THMs concentration, the regression equation indicates that, on the average, for 90 min increases in contact time caused decreases of THM4 by about (90.8%).




Figure 10: Variation of TTHMs Concentration with Contact Time

The variation of individual compound with the contact time is shown in Figure 11. The concentration of chloroform, bromodichloromethane, dibromochloromethane and bromoform decreased with increasing detention time, a production curve almost parallel to those TTHMs significantly lower with increasing contact time.




Figure 11: Variation of TTHMs Compound with Contact Time

Figure 12 represents the percent reduction in TTHMs concentration along 2 hour detention time. This Figure shows that the largest percent removal of TTHMs is (94.43%) at (120min) of detention time. Thus, we conclude that the reduction in THMs concentration is proportional with time in the form ($Y=0.0057X+0.29$) with coefficient of determination ($R^2=0.88$).




Figure 12: Percent Reduction in TTHMs with the Contact Time

When change ClO_2 contact time the average weight fraction of bromoform compound ranged from 2.97% to 6.4%, while the average weight fraction of CHBr_3 was 27.72% to 35.8%, for CHBr_2Cl was 30% to 37.65%, and for CHClBr_2 was 23.58% to 30.17%) as shown in Figure 13.




Figure 13: Weight Fraction of Individual THM4 to Total THM4

Figure 14 shows the relation between detention time and percent reduction of four compounds (chloroform, bromodichloromethane, dichlorobromomethane, and bromoform), as shown the percent removal of four compound of THMs increases with increasing contact time and higher percent removal of THMs compounds is 97.13%, 96%, 95.09%, 87.20% at 120 min for bromoform, dichlorobromomethane, bromodichloromethane, and chloroform respectively.




Figure 14: Percent Reduction of THMs Compounds with the Detention Time

TOTAL ORGANIC CARBON CONCENTRATIONS IN TREATED WATER

Figure 15 represents a plot of variations in TOC with various chlorine dioxide doses. As can be seen, the concentrations varied little throughout the entire study period. Additionally, TOC concentrations remaining after each run were not significantly different from one another, but it is decrease with increase ClO_2 dose.

Results show that the trend between total organic carbon concentration and chlorine Dioxide Dose began to decreases significantly with the increasing ClO_2 dose. Thus, we conclude that the TOC concentration is proportional with ClO_2 dose with coefficient of determination ($R^2 = 0.93$) in the form ($Y = -0.25X + 1.438$).




Figure 15: Variation of Total Organic Carbon with ClO_2 Dose

Figure 16 shows the relation between TTHMs and TOC concentration. The correlations are positive between them; the result shows that the TTHMs concentration increases significantly with the increasing total organic carbon. So, we conclude that the THMs concentration is proportional with TOC in the form ($Y=0.046X+0.015$) with coefficient of determination ($R^2=0.96$).




Figure 16: Variations in TTHMs with TOC

EFFECT OF TURBIDITY ON TTHMS FORMATION

Turbidity of filtered water ranges between (4 to 7.36) NTU. Lower value when water pre oxidation with (2mg/L) ClO_2 dose while the higher value for (0.5mg/L) ClO_2 . Figure 17 shows that increasing in turbidity lead to increase TTHMs concentration, the correlations are positive between them.

Results show that the TTHMs concentration increases significantly with the increasing turbidity. So, we conclude that the THMs concentration is proportional with turbidity, and the correlations are positive between them, in the form ($Y=0.0057X+0.037$) with coefficient of determination ($R^2=0.85$).




Figure 17: Variation of Turbidity with Trihalomethane

Figure 18 shows the variation of four THMs compound with turbidity. As show the result of CF, BDCM, BF, and DCBM variation began slightly decrease with increasing turbidity.




Figure 18: Variation of THMs Compounds with Turbidity

EFFECT OF PH ON TRIHALOMETHANES

Figure 19 shows the variation of TTHMs with pH in all runs. It has been found that the degree of inactivation by chlorine dioxide increases as pH increases leads to increases of THMs formation. This finding is agreed with the results adapted by (Bernarde et al, 1967; Letterman 1999; AWWA, 1982).

pH effects on THMs formation mechanisms and thus speciation, in which increasing pH lead to increase hydrolysis reactions that many DBPs undergo to form THMs such as trihalopropanones, trihaloacetonitriles, trihaloacetaldes, trihalonitromethanes and trihaloacetic acids (Nikolaou, 2004). The relation between pH and THMs formation is in the form ($Y=0.018X+0.071$) with coefficient of determination R^2 equal to (0.973).




Figure 19: Variation of pH with TTHMs

EFFECT OF TEMPERATURE ON TRIHALOMETHANES

The temperatures ranged in all runs throughout the study period between (24-30) °C. The variation of temperature with THMs is shown in Figure (20). The results show slight variation in TTHMs with temperature, but it is very recognized that TTHMs increase as temperature increase, in the form ($Y=0.0032X-0.0154$) with coefficient of determination ($R^2=0.823$), because the disinfection efficiency of chlorine dioxide decreases as temperature decreases. This finding is agreed with the results adapted by (Bernarde, et al., 1967).




Figure 20: The Variation of Temperature with TTHM

CONCLUSIONS

Based on the objectives of this study the following conclusions and are presented:-

- The results showed that the THM formation decrease with increasing chlorine dioxide dose, concentrations of THMs that developed in water pretreated with 2 mg/L ClO_2 were significantly lower when compared to those in water pretreated with the other dose of ClO_2 .
- The ability of ClO_2 to lowering THMs formation was very good. Increasing dose of ClO_2 from 1 mg/L to 2 mg/L will caused (24%) reduction of THMs concentrations.
- Chlorine dioxide showed a good potential as an alternative disinfectant and oxidant produced lower THMs concentrations but only at doses higher than (1 mg/L). The optimal dose of ClO_2 produced high percent reduction of THM formation (64.33 %) is (2) mg/L.
- The THMs concentration is proportional with time in the form ($Y = -0.0009X + 0.1088$) with coefficient of determination $R^2 = 0.88$, increasing of chlorine dioxide contact time from (30 to 120 min) will caused (90.85%) reduction in THMs formation, and higher percent removal of THMs compounds is 97.13% , 96%, 95.09%, 87.20% at 120 min for bromoform, dichlorobromomethane, bromodichloromethane, and chloroform respectively.
- Chloroform concentration is proportional with ClO_2 dose in the form ($Y = 5.3961X + 52.04$) with coefficient of determination ($R^2 = 0.0407$), and the correlation was negative under the dose 1mg/L.
- Correlations are positive between bromodichloromethane and dichlorobromomethane percent reduction and chlorine dioxide dose and slightly decrease in bromoform concentrations with increasing chlorine dioxide dose.
- All results indicate that the treated water contain mainly the brominate THM species (CHClBr_2 and CHBrCl_2). The average weight fraction of chloroform (CHCl_3) compound ranged from 7.16% to 21.46%, while the average weight fraction of the rest brominated (CHBr_2Cl , and CHBrCl_2) species ranged from about 34.97% to 43.8%, 30.67% to 34.54% respectively.
- Results show that the trend between total organic carbon concentration and chlorine dioxide dose began to decreases significantly with the increasing ClO_2 dose, TTHMs concentration increases significantly with the increasing total organic carbon, and turbidity.
- The relation between pH and THMs formation is in the form ($Y = 0.018X + 0.071$) with coefficient of determination R^2 equal to (0.973), and found that the degree of inactivation by chlorine dioxide increases as pH increases.
- The results show slight variation in TTHMs with temperature, in the form ($Y = 0.0032X - 0.0154$) with coefficient of determination ($R^2 = 0.823$).
- Comparison of total THMs measurements with the USEPA Stage I and stage II limits. Shows that none of the THM levels exceeded the stage I limit, while exceeded the stage II limit in all sampling measurements.
- Dominant TTHMs species during all runs is DCBM, it consists (39.2%) of trihalomethanes.

REFERENCES

1. **Abedin, Azizi1; Kadri, Berisha1; Selim, Jusufi2; Shefqet, Rashani1; 2010**, "Formation of Disinfection By-products (DBPs) and Strategies to Reduce their Concentration in the Water Treatment Plants in Përlepnica and Velekince - Gjilan", Faculty of Mines and Metallurgy, Department of Technology, University of Prishtina.

2. **AWWA, 1982**, "*Treatment Techniques for Controlling Trihalomethanes in Drinking Water*", Denver, CO: AWWA.
3. **Bernarde, M.A; W.B., Snow; and V.P. Olivieri, 1967**, "*Chlorine Dioxide Disinfection Temperature Effects*", J. Appl. Bacteriol. 30(1):159.
4. **Griese, Mark, 1991**, "*Pilot Plant Optimization of the Chlorine Dioxide Treatment Process for Disinfection By-Product Reduction*", Conference Proceedings, AWWA Annual Conference, Philadelphia, PA.
5. **Hulsey, R.A.; J.J. Neemann; S.N. Foellmi; and R.C. Hoehn, 2000**, "*Chlorine Dioxide for DBP Control and Filter Enhancement*", Conference Proceedings, AWWA Water Quality Technology Conference, Salt Lake City, Utah
6. **Krasner S.W; Scilimenti, M. J.; Means, E. G.; Symons, J. M.; and Simms, L., 1992**, "*The Impact of chlorine dose, residual, chlorine, bromide, and organic carbon on trihalomethane speciation*", AWWA Water Quality Technology Conference.
7. **Letterman, Raymond, 1999**, "*Water Quality and Treatment: A Handbook of Community Water Supplies* ", New York, NY: McGraw-Hill, Inc.
8. **McLain, J.L.; A. Obolenski; and H.M. Shukairy; 2002**, "*Disinfection By-Product Speciation*", AWWA, 141-168.
9. **Stevens, A.A.; L.A., Moore; and R.J, Mifflner, 1989**, "*Formation and Control of Non trihalomethane Disinfection By-products*", J.AWWA, 81:8:54-60.
10. **Babcock, D. B.; and Singer, P. C., 1979**, "Chlorination and coagulation of humic and fulvic acids", Proc. 97th Ann. AWWA Conf. Paper 16-6.
11. **Fawell, J. K.; and Hunt, S., 1988**, "Chloroform and trihalomethanes. In Environmental Toxicology: Organic Pollutants" Chapters 1 and 2, pp. 15-36. Ellis Horworth Ltd, Chichester, West Sussex
12. **Nikolaou, A. D; Lekkas, T.D; and Golfinopoulos, S.K, 2004**, "Kinetics of the formation and decomposition of chlorination by-products in surface waters", Chemical engineering Journal, 100, 139.
13. **Kavanough, M. C; Trussell, A. R.; Cromer, J.; and Trussell, R. R.; 1980**, "An empirical Kinetic model of THM formation: applications to meet the proposed THM standards", J. Am. Water Works Assoc. 72, 578-582.
14. **Klein, S., 1990**, "Bildung von organohalogenverbindungen bei der wasserchlorung", Z. Gesamte Hyg. 36, 32-535.
15. **EEC, 1980**, "Directiva del Consejo de 15 de Julio de 1980 relativa a la calidad de las aguas destinadas al consumo humano 80/778/CEE". Diario Oficial de las Comunidades Europeas, no. L229/11.
16. **Peters, C. J.; Young, R. J.; and Perry, R., 1980**, "Factors influencing the formation of haloforms in the chlorination of humic materials", Envir. Sci. Technol. 14, 1391-1395.
17. **Trussell, R. R.; and Umphres, M. D., 1978**, "The formation of trihalomethanes", J. Am. Water Works Assoc. 70, 604-612. A. A., Slocum C. J., Seeger D. R. and
18. **Onodera, S.; Nishikawa, T.; and Suzuki, S., 1987**, "Chemical changes in chlorinated water.

- XIV Characterization and determination of halogenated organics formed during chlorination of water from the Tama River”, J. Chromatogr. 409, 259-270.
19. **Rook, J. J., 1974**, "Formation of Haloforms during Chlorination of Natural Waters", Water Treatment Exam, 23:2:234-243.
 20. **Simmon, V. F.; and Tardiff, R. G., 1978**, "The mutagenic activity of halogenated compounds found in chlorinated drinking water", In Water Chlorination: Environmental Impact and Health Effects (Edited by Jolley R. L., Gorcher H. and Hamilton D. H.), Vol. 2, pp. 417-432. Ann Arbor Science, Ann Arbor, MI.
 21. **Sandier, R., 1977**, "Untersuchungensurbuildung von organischenchlorrerbindungenbei der chlorunghuminsaurehaltigerwasser", Thesis, Engler-Bunte Institute, University of Karlsruhe.
 22. **Trehy, M. L.; and Bieber, T. S., 1980**, "Effect of commonly used water treatment processes on the formation of THM", Proc. AWWA Ann. Conf. 1980, pp. 125-138.
 23. **USEPA Drinking Water, 1988**, "Substitution of contaminants and drinking water priority list of additional substances which may require regulation under the Safe Drinking Water Act", Fed. Reg. 53(14), 1892.
 24. **WHO, 1992**, "Revision of the WHO guidelines for drinking water quality. Report of the final task group meeting", Geneva (limited distribution document).

